Subject Index

Acremonium lolii: see Nematodes, plantparasitic, 403

Aerial spraying: see Permethrin, 357 Agaricus bisporus: see Biocontrol, 271

Agelaius phoeniceus: Oil concentration differences among sunflower achenes and feeding preferences of redwinged blackbirds, 294

Agromyza sp.: see Bionomics, 70

Alternaria macrospora: Suppression of Alternaria leaf spot in Pima cotton by systemic fungicides, 381

Animal grazing: see Growth regulators, plant, 178

Antagonists: see Biocontrol, 172

Antifeedants: see Semiochemicals, 145 Antitrogus consanguineus: see Sugar-cane whitegrub, 205

Application method: see Weed control. 254

Application methods: Distribution of dve deposits from a small-plot track sprayer on biological and artificial targets, 129

Application methods: see Insecticides, 111

Application schedules: Calendar and monitored insecticide application for the control of cowpea pests, 363

Bacillus thuringiensis: see Traps, pheromone, 229

Banana weevil borer: see Resistance, 215 Barley powdery mildew: see Fungicide resistance, 445

Bemisia tabaci: see Gossypium spp., 461 Bifenthrin: see Whiteflies, 106

Bioassay, knockdown: see pisorum, 53

Bioassay: see Liriomyza spp., 509

Biocontrol: Biocontrol of plant diseases: the approach for tomorrow, 95

Biocontrol: Biological control fusarium diseases by fluorescent Pseudomonas and non-pathogenic Fusarium, 279

Biocontrol: Biological control of postharvest diseases of fruits and vegetables: alternatives to synthetic fungicides, 172

Biocontrol: Development and application of a biocontrol system for bacterial blotch of the cultivated mushroom, 271

Biocontrol: Diversification of biological control strategies in agriculture, 243

Biocontrol: Diversity in biological control. 85

Biocontrol: Seed treatments for biological control of plant disease, 166

Biocontrol formulations: Biocontrol of cotton damping-off caused by

Rhizoctonia solani in the field with formulations of Trichoderma spp. and Gliocladium virens, 396

Bionomics: Bionomics of the wheat leafminer Agromyza sp. (Diptera: Agromyzidae) in Central Saudi Arabia, 70

Broccoli cultivars: Cultivar preferences of lepidopterous pests of broccoli,

Bruchus pisorum: Bruchus pisorum L. (Coleoptera, Bruchidae) control by a knockdown pyrethroid in field peas,

Cavity spot: see Seed treatment, 117 Cercospora sojina: see Glycine max, 465 Chara: Control of Chara with oxadiazon

and copper sulphate in waterlogged rice fields in India, 371 Colomerus vitis: see Grape vine, 310

Control strategy: see Model, 209 Control, chemical: New developments in

chemical control of white root disease of Hevea brasiliensis in Africa, 234

Control, chemical: see Nematodes, 121 Cotton: see Electrostatics, 354 Cotton: see Insecticides, 111

Cotton: see Integrated pest management,

Cowpea pests: see Application schedules, 363

Cowpea: Laboratory and screenhouse assays for evaluating cowpea resistance to the legume pod borer, 48

Cowpea: see Nematodes, 121 Crop damage: Disease-induced losses in winter wheat in England and Wales 1985-1989, 504

Crop damage: see Empoasca fabae, 416 Crop damage: see Preference, 455

Crop damage: see Sugar-cane borer, 57 Crop damage: see Sugar-cane whitegrub,

Crop yield: see Glycine max, 465 Crop yield: see Rust, leaf, 222 Cropping sequence: see Weed control, 28 Cropping systems: see Whiteflies, 305 Cucumber: see Resistance, induced, 265

Cucumber: see Seedling diseases, 23

Degradation: Degradation of pesticides by micro-organisms and the potential for genetic manipulation, 334

Drop size: Pesticide droplet size and efficacy: a series of trials on grasshoppers, 136

Drop size: Variations in insecticide dose received by settled locusts: a computer model for ultra-low-volume spraying, 183

ELISA: see Rice tungro virus, 195

Economics: Decision support system for economic analysis of grasshopper treatment operations in the African Sahel 485

Economics: see Take-all, 391

Electrostatics: Studies on the performance of the 'Electrodyn' on cotton in eastern Tanzania, 354

Empoasca fabae: Validation of a model simulating the feeding effects of the potato leafhopper (Empoasca fabae) on potato growth, 416

Empoasca lybica: see Gossypium spp., 461 Ervsiphe graminis f.sp. hordei: see Fungicide resistance, 445

Fentin acetate: Effect of fentin acetate on wet-seeded rice, Pistia stratiotes and Azolla pinnata, 45

Fluid drilling: see Seedling diseases, 23 Foliar-applied urea: Use of foliar-applied urea to inhibit the development of Septoria nodorum in spring wheat, 260

Forecasting: Predicting the survival of immigrant insect pests in new environments, 331

Forecasting: see Slug, 413

Forestry: see Permethrin, 357 Formulation: see Persistence, 41

Fungicide: see Alternaria macrospora, 381 Fungicide mixtures: see Pythium aphanidermatum, 386

Fungicide resistance: see Pseudocercosporella herpotrichoides, 479

Fungicide resistance: Sensitivity of Erysiphe graminis f.sp. hordei to morpholine and piperidine fungicides,

Fungicides: see Rhizoctonia root rot, 473 Fusarium oxysporum: see Biocontrol, 279

Genetic manipulation: see Biocontrol, 85 Genetic manipulation: see Biocontrol, 95 Genetic manipulation: see Degradation,

Glycine max: Detached shoot technique to evaluate the reaction of soybean cultivars to Sclerotium rolfsii, 325

Glycine max: Yield losses in soybeans from frogeye leaf spot caused by Cercospora sojina, 465

Gossypium barbadense: see Alternaria macrospora, 381

Gossypium hirsutum: see Biocontrol formulations, 396

Gossypium spp.: Cotton leaf trichomes and populations of Empoasca lybica and Bemisia tabaci, 461

Grain protectants: see Insect pests, 287 Grape vine: A pruning technique for

Subject Index

Acremonium lolii: see Nematodes, plantparasitic, 403

Aerial spraying: see Permethrin, 357 Agaricus bisporus: see Biocontrol, 271

Agelaius phoeniceus: Oil concentration differences among sunflower achenes and feeding preferences of redwinged blackbirds, 294

Agromyza sp.: see Bionomics, 70

Alternaria macrospora: Suppression of Alternaria leaf spot in Pima cotton by systemic fungicides, 381

Animal grazing: see Growth regulators, plant, 178

Antagonists: see Biocontrol, 172

Antifeedants: see Semiochemicals, 145 Antitrogus consanguineus: see Sugar-cane whitegrub, 205

Application method: see Weed control. 254

Application methods: Distribution of dve deposits from a small-plot track sprayer on biological and artificial targets, 129

Application methods: see Insecticides, 111

Application schedules: Calendar and monitored insecticide application for the control of cowpea pests, 363

Bacillus thuringiensis: see Traps, pheromone, 229

Banana weevil borer: see Resistance, 215 Barley powdery mildew: see Fungicide resistance, 445

Bemisia tabaci: see Gossypium spp., 461 Bifenthrin: see Whiteflies, 106

Bioassay, knockdown: see pisorum, 53

Bioassay: see Liriomyza spp., 509

Biocontrol: Biocontrol of plant diseases: the approach for tomorrow, 95

Biocontrol: Biological control fusarium diseases by fluorescent Pseudomonas and non-pathogenic Fusarium, 279

Biocontrol: Biological control of postharvest diseases of fruits and vegetables: alternatives to synthetic fungicides, 172

Biocontrol: Development and application of a biocontrol system for bacterial blotch of the cultivated mushroom, 271

Biocontrol: Diversification of biological control strategies in agriculture, 243

Biocontrol: Diversity in biological control. 85

Biocontrol: Seed treatments for biological control of plant disease, 166

Biocontrol formulations: Biocontrol of cotton damping-off caused by

Rhizoctonia solani in the field with formulations of Trichoderma spp. and Gliocladium virens, 396

Bionomics: Bionomics of the wheat leafminer Agromyza sp. (Diptera: Agromyzidae) in Central Saudi Arabia, 70

Broccoli cultivars: Cultivar preferences of lepidopterous pests of broccoli,

Bruchus pisorum: Bruchus pisorum L. (Coleoptera, Bruchidae) control by a knockdown pyrethroid in field peas,

Cavity spot: see Seed treatment, 117 Cercospora sojina: see Glycine max, 465 Chara: Control of Chara with oxadiazon

and copper sulphate in waterlogged rice fields in India, 371 Colomerus vitis: see Grape vine, 310

Control strategy: see Model, 209 Control, chemical: New developments in

chemical control of white root disease of Hevea brasiliensis in Africa, 234

Control, chemical: see Nematodes, 121 Cotton: see Electrostatics, 354 Cotton: see Insecticides, 111

Cotton: see Integrated pest management,

Cowpea pests: see Application schedules, 363

Cowpea: Laboratory and screenhouse assays for evaluating cowpea resistance to the legume pod borer, 48

Cowpea: see Nematodes, 121 Crop damage: Disease-induced losses in winter wheat in England and Wales 1985-1989, 504

Crop damage: see Empoasca fabae, 416 Crop damage: see Preference, 455

Crop damage: see Sugar-cane borer, 57 Crop damage: see Sugar-cane whitegrub,

Crop yield: see Glycine max, 465 Crop yield: see Rust, leaf, 222 Cropping sequence: see Weed control, 28 Cropping systems: see Whiteflies, 305 Cucumber: see Resistance, induced, 265

Cucumber: see Seedling diseases, 23

Degradation: Degradation of pesticides by micro-organisms and the potential for genetic manipulation, 334

Drop size: Pesticide droplet size and efficacy: a series of trials on grasshoppers, 136

Drop size: Variations in insecticide dose received by settled locusts: a computer model for ultra-low-volume spraying, 183

ELISA: see Rice tungro virus, 195

Economics: Decision support system for economic analysis of grasshopper treatment operations in the African Sahel 485

Economics: see Take-all, 391

Electrostatics: Studies on the performance of the 'Electrodyn' on cotton in eastern Tanzania, 354

Empoasca fabae: Validation of a model simulating the feeding effects of the potato leafhopper (Empoasca fabae) on potato growth, 416

Empoasca lybica: see Gossypium spp., 461 Ervsiphe graminis f.sp. hordei: see Fungicide resistance, 445

Fentin acetate: Effect of fentin acetate on wet-seeded rice, Pistia stratiotes and Azolla pinnata, 45

Fluid drilling: see Seedling diseases, 23 Foliar-applied urea: Use of foliar-applied urea to inhibit the development of Septoria nodorum in spring wheat, 260

Forecasting: Predicting the survival of immigrant insect pests in new environments, 331

Forecasting: see Slug, 413

Forestry: see Permethrin, 357 Formulation: see Persistence, 41

Fungicide: see Alternaria macrospora, 381 Fungicide mixtures: see Pythium aphanidermatum, 386

Fungicide resistance: see Pseudocercosporella herpotrichoides, 479

Fungicide resistance: Sensitivity of Erysiphe graminis f.sp. hordei to morpholine and piperidine fungicides,

Fungicides: see Rhizoctonia root rot, 473 Fusarium oxysporum: see Biocontrol, 279

Genetic manipulation: see Biocontrol, 85 Genetic manipulation: see Biocontrol, 95 Genetic manipulation: see Degradation,

Glycine max: Detached shoot technique to evaluate the reaction of soybean cultivars to Sclerotium rolfsii, 325

Glycine max: Yield losses in soybeans from frogeye leaf spot caused by Cercospora sojina, 465

Gossypium barbadense: see Alternaria macrospora, 381

Gossypium hirsutum: see Biocontrol formulations, 396

Gossypium spp.: Cotton leaf trichomes and populations of Empoasca lybica and Bemisia tabaci, 461

Grain protectants: see Insect pests, 287 Grape vine: A pruning technique for

saving vineyards severely infested by the grape vine bud mite Colomerus vitis (Pagenstecher) (Eriophyidae).

Grape vine: see Parasitoids, 501 Grasshoppers: see Drop size, 136 Grasshoppers: see Drop size, 183

Grasshopper, Senegalese: see Economics,

Growth regulators, insect: see Semiochemicals, 145

Growth regulators, plant: Regulating sward growth and animal output with mefluidide, 178

Glycine max: Leaf-feeding resistance to six Australian noctuids in sovbean. 320

Harrisina brillians: see Parasitoids, 501

Herbicide: see Chara, 371

Herbicide: Control of Sinapis arvensis in safflower with post-emergence herbicides, 74

Herbicide: see Weed control, 254

Herbicides: Effectiveness of selective herbicides for the control of annual weed species in lathyrus (Lathyrus sativus L.), 496

Hevea brasiliensis: see Control, chemical,

Insect control: see Economics, 485 Insecticidal control: see Application schedules, 363

Insecticide resistance: see Liriomyza spp., 500

Insecticide resistance: see Whiteflies, 106 Insecticides: Effect of swath width, time of application and height on the efficacy of very-low-volume (VLV) water-based insecticides on cotton in Malawi, 111

Insect pests: Assessment and control of losses caused by insect pests in small farmers' stores in Zimbabwe, 287

Insect pests: see Broccoli cultivars, 199 Insect pests: see Forecasting, 331

Insect pests: Insect pests of spring wheat in Central Saudi Arabia, 65

Integrated control: see Parasitic weeds, 6 Integrated management: Integrated management of rice tungro disease in South Sulawesi, Indonesia, 34

Integrated pest management: Prospects and status for development of novel chemicals for IPM in cotton, 347

Jojoba: Sucking insects damaging jojoba, Simmondsia chinensis (Link) Schneider, and their natural enemies, in the North Central and Central regions of Chile, 469

Lathyrus sativus: see Herbicides, 496 Legume seeds: see Oil treatments, 375 Legume seeds: see Stored product, 315 Liriomyza spp.: Responses of the alien leaf miners Liriomyza trifolii and

Liriomyza huidobrensis (Diptera: Agromyzidae) to some pesticides scheduled for their control in the UK.

Maruca testulalis: see Cowpea, 48 Mefluidide: see Growth regulators, plant,

Meloidogyne incognita: see Tomato, 62 Metalaxyl: see Seed treatment, 117 Meteorological: see Slug. 413

Methiocarb: see Slug. 408 Model: see Empoasca fabae, 416

Model: Evaluation by computer simulation of strategies to time metalaxyl applications for improved control of potato late blight, 209

Models, logistic: Deterministic and stochastic logistic models for describing increase of plant diseases, 155

Models, mathematical: Mathematical models in weed management, 432

Natural enemies: see Biocontrol, 243 Natural enemies: see Jojoba, 469

Nematodes: Chemical control of the nematode pests of cowpea. Vigna unguiculata (L.) Walp, 121

Nematodes, plant-parasitic: Effects on plant-parasitic nematodes of infection of perennial ryegrass, Lolium perenne, by the endophytic fungus, Acremonium lolii, 403

Nephotettix spp: see Integrated management, 34

Nilaparvata lugens: Impact of resistance mechanisms in rice against the brown planthopper, Nilaparvata lugens Stal (Homoptera: Delphacidae), 125

Noctuid larvae: see Glycine max, 320

Oil treatments: Protection of stored legume seeds against attack by storage fungi and weevils: mechanism of action of lipoidal and oil seed treatments, 375

Oil treatments: see Stored product, 215 Organophosphorus insecticides: Resistance, 215

Parasitic weeds: Protection of crops against parasitic weeds, 6

Parasitoids: Parasitoids of the grapeleaf skeletonizer, Harrisina Barnes and McDonnough (Lepidoptera: Zygaenidae) in northwestern Mexico, 501

Permethrin: Airborne permethrin and off-target deposits from an aerial ultra-low-volume silvicultural spray, 357

Permethrin: see Persistence, 41

Persistence: Influence of formulation and spray droplet size upon the persistence of permethrin deposits on brussels sprout leaves, 41

Pest damage: see Jojoba, 469

Pesticides: see Degradation, 334 Pesticides: see Drop size, 136

Pesticides: see Integrated pest management 347

Phytophthora infestans: see Model, 209 Phytotoxicity: see Fentin acetate, 45 Phytotoxicity: see Herbicides, 496 Pisum sativum: see Bruchus pisorum, 53

Plant diseases: see Biocontrol 85 Plant diseases: see Biocontrol, 95

Plant diseases: see Biocontrol, 166 Plant diseases: see Crop damage, 504 Plant diseases: see Models, logistic, 155 Plant diseases: see Resistance, induced.

265 Population dynamics: see Forecasting.

Population dynamics: see Slug, 408 Population dynamics: see Whiteflies, 305 Post-harvest diseases: see Biocontrol, 172 Preference: see Agelaius phoeniceus, 294 Preference: see Broccoli cultivars, 199

Preference: Sucrose avoidance by American robins (Turdus migratorius): implications for control of bird damage in fruit crops, 455

Pruning: see Grape vine, 310

Pseudocercosporella herpotrichoides: Effects of fungicides on evespot in five successive crops of winter wheat in plots with different initial or fungicide-selected populations of Pseudocercosporella herpotrichoides,

Pseudomonas, fluorescent: see Biocontrol,

Pseudomonas, fluorescent: see Biocontrol. 279

Pterohelaeus: see Seed treatment, 293 Pythium aphanidermatum: Synergistic and antagonistic interactions of fungicides against Pythium aphaniderm-

atum on perennial ryegrass, 386

Resistance: Organophosphorus insecticide resistance and its management in the banana weevil borer, Cosmopolites sordidus (Germar) (Coleoptera: Curculionidae), in Australia, 215

Resistance breeding: see Biocontrol, 243 Resistance, induced: Oxalate and phosphates induce systemic resistance against diseases caused by fungi, bacteria and viruses in cucumber, 265

Resistance mechanisms: see Nilaparvata lugens, 125

Resistance screening: see Cowpea, 48 Resistance screening: see Glycine max, 320

Resistance screening: see Glycine max, 325

Resistance screening: see Rice tungro virus, 195

Rhizoctonia root rot: Evaluation of infurrow fungicide treatments to control rhizoctonia root rot of wheat, 473

Rhizoctonia solani: see Biocontrol formulations, 396

Rice tungro virus: Efficient method of determining tungro virus resistance

- in rice (Oryza sativa L.), 195
- Rice tungro virus: see Integrated management. 34
- Rice: see Chara, 371
- Rice varieties: see Nilaparvata lugens, 125
- Rust, leaf: Influence of leaf position on yield loss caused by wheat leaf rust in single tillers, 222
- Ryegrass: see Pythium aphanidermatum, 386
- Ryegrass: see Nematodes, plantparasitic, 403
- Safflower: see Herbicide, 74
- Saudi Arabia: see Bionomics, 70
- Saudi Arabia: see Insect pests, 65
- Sclerotium rolfsii: see Glycine max, 325
 Seed treatment: Effect of film-coat layering of metalaxyl on the germination
 of carrot seeds, their emergence and
- the control of cavity spot, 117
 Seed treatment: Soil-insecticide bioassays
 and seed treatments for soil-insect
 control in Central Queensland, Australia, 293
- Seed treatment: see Biocontrol, 166
- Seedling diseases: Stand and yield of cucumber seeded with gel and fungicides in various tillage systems, 23
- Semiochemicals: Laboratory evaluation of pest management strategies combining antifeedants with insect growth regulator insecticides, 145
- Septoria nodorum: see Foliar-applied urea, 260
- Sinapis arvensis: see Herbicide, 74
- Slug: Comparison between defined-area slug traps and other methods of trapping slugs in cereal fields, 152
- Slug: Development of a forecast of slug activity: models to relate slug activity to meteorological conditions, 413
- Slug: Slug population changes following

- molluscicide treatment in relation to distance from edge of treated area,
- Soil insecticides: see Seed treatment, 293 Soils, arable: see Slug, 152
- Spodoptera exempta: see Traps, pheromone, 229
- Spray deposit: see Application methods,
- Spray deposit: see Drop size, 183
- Stalk rot: see Sugar-cane borer, 57 Stored product: Efficacy of oil treatments of legume seeds for control of Aspergillus and Zabrotes, 315
- Stored product: see Insect pests, 287 Stored product: see Oil treatments, 375
- Sugar-cane borer: Effects of *Diatraea sac*charalis (F.) (Lepidoptera: Pyralidae) damage and stalk rot fungi on sugarcane yield in Louisiana, 57
- Sugar-cane whitegrub: Effect of Antitrogus consanguineus (Blackburn) (Coleoptera: Scarabacidae) infestations on sugar-cane yield in Australia, 205
- Sunflower: see Agelaius phoeniceus, 294
- Take-all: Economic significance of takeall on winter wheat in the Pacific Northwest USA, 391
- Tillage: see Weed control, 28
- Tomato: Reaction of three tomato cultivars to two vascular diseases in presence of the root-knot nematode, Meloidogyne incognita Race 1, 62
- Track spacing: see Electrostatics, 354
- Track sprayer: see Application methods, 129
- Trap: see Slug, 152
- Traps, pheromone: Monitoring secondary outbreaks of the African armyworm in Kenya using pheromone traps for timing of *Bacillus thuringien*sis application, 229

- Turdus migratorius: see Preference, 455
- Vascular wilts: see Tomato, 62
- Weed control: Influence of weed-control practices in the first crop on the tillage requirements for the succeeding crops in an upland rice-maize-cowpea cropping sequence, 28
- Weed control: see Models, mathematical, 432
- Weed control: see Parasitic weeds, 6
 Weed control: Potential for automatic
 weed detection and selective herbicide application, 254
- Weed-crop interactions: see Models, mathematical, 432
- Wet-seeded rice: see Fentin acetate, 45 Wheat: see Rhizoctonia root rot, 473
- Wheat, spring: see Foliar-applied urea,
- Wheat, spring: see Insect pests, 65
- Wheat, spring: see Rust, leaf, 222
- Wheat, winter: see Crop damage, 504 Wheat, winter: see Pseudocercosporella
- herpotrichoides, 479
- Wheat, winter: see Take-all, 391
- White root disease: see Control, chemical, 234
- Whiteflies: Efficacy of bifenthrin against pyrethroid-resistant and susceptible populations of glasshouse whitefly in bioassays and using three spray application methods, 106
- Whiteflies: Survivorship of the cassava whiteflies Aleurotrachelus socialis and Trialeurodes variabilis (Homoptera: Aleyrodidae) under different cropping systems in Colombia, 305
- Zucchini yellow mosaic virus: see Models, logistic, 155

Author Index

ADAMS, A. J., LINDQUIST, R. K., ADAMS, I. H. H. AND HALL, F. R. Efficacy of bifenthrin against pyrethroid-resistant and -susceptible populations of glasshouse whitefly in bioassays and using three spray application methods, 106

ADAMS, I. H. H. see ADAMS, A. J., 106
AFUN, J. V. K., JACKAI, L. E. N. AND
HODGSON, C. J. Calendar and monitored insecticide application for the control of cowpea pests, 363

AKEM, C. N. see DASHIELL, K. E., 465 AKEM, C. N. AND DASHIELL, K. E. Detached shoot technique to evaluate the reaction of soybean cultivars to Sclerotium rolfsii, 325

ALABOUVETTE, C. see LEMANCEAU, P., 279
ALLSOPP, P. G., BULL, R. M. AND
MCGILL, N. G. Effect of Antirrogus
consanguineus (Blackburn) (Coleoptera: Scarabaeidae) infestations on
sugar-cane yield in Australia, 205

ALTIERI, M. A. see GOLD, C. S., 305 Araya, J. E. see Quiroga, D. 469 Arretz, P., see Quiroga, D., 469

BABATOLA, J. O. AND OMOTADE, M. A. Chemical control of the nematode pests of cowpea, *Vigna unguiculata* (L.) Walp, 121

BAILEY, P. see HORNE, J., 53

BAKER, R. Diversity in biological control, 85

BATEMAN, G. L. AND FITT, B. D. L. Effects of fungicides on eyespot in five successive crops of winter wheat in plots with different initial or fungicide-selected populations of Pseudocercosporella herpotrichoides, 470

BELLOTTI, A. C. see GOLD, C. S., 305 BILES, C. L. see WILSON, C. L., 172

BRIER, H. B. AND ROGERS, D. J. Leaffeeding resistance to six Australian noctuids in soybean, 320

Brown, J. K. M., Slater, S. E. and See, K. A. Sensitivity of *Erysiphe graminis* f.sp. *hordei* to morpholine and piperdine fungicides, 455

BROWNBRIDGE, M. see BROZA, M., 229
BROZA, M., BROWNBRIDGE, M. AND
SNEH, B. Monitoring secondary outbreaks of the African armyworm in
Kenya using pheromone traps for
timing of Bacillus thuringiensis application, 229

BRUGGER, K. E. AND NELMS, C. O. Sucrose avoidance by American robins (*Turdus migratorius*): implica-

tions for control of bird damage in fruit crops, 455

Bull, R. M. see Allsopp, P. G., 205
Butler, G. D., Wilson, F. D. and Fish-Ler, G. Cotton leaf trichomes and populations of *Empoasca lybica* and *Bemisia tabaci*, 461

BUTLER, R. C. see GLEN, D. M., 408

CABUNAGAN, R. C. see SAMA, S., 34
CHALUTZ, E. see WILSON, C. L., 172
CHELLIAH, S. see SENGUTTUVAN, T., 125
CHINDO, P. S., KHAN, F. A. AND ERINLE,
I. D. Reaction of three tomato cultivars to two vascular diseases in presence of the root-knot nematode,
Meloidogyne incognita Race 1, 62

CLEAR HILL, B. H. see HAGGAR, R. J., 178
CLEMENTS, R. O. AND MURRAY, P. J.
Comparison between defined-area
slug traps and other methods of trapping slugs in cereal fields, 152

COLLINS, P. J., TREVERROW, N. L. AND LAMBKIN, T. M. Organophosphorus insecticide resistance and its management in the banana weevil borer, Cosmopolites sordidus (Germar) (Coleoptera: Curculionidae), in Australia, 215

СООК, R. J. see FOLWELL, R. J., 391 СООК, R., LEWIS, G. C. AND MIZEN, K. A. Effects on plant-parasitic nematodes of infection of perennial ryegrass, Lolium perenne, by the endophytic

fungus, Acremonium Iolii, 403 Cook, R. J., Polley, R. W. and Thomas, M. R. Disease-induced losses in winter wheat in England and Wales 1985–1989, 504

COOP, L. B., CROFT, B. A., MURPHY, C. F. AND MILLER, S. F. Decision support system for economic analysis of grasshopper treatment operations in the African Sahel, 485

COTTERILL, P. J. Evaluation of in-furrow fungicide treatments to control rhizoctonia root rot of wheat, 473

COUCH, H. B. AND SMITH, B. D. Synergistic and antagonistic interactions of fungicides against *Pythium* aphanidermatum on perennial ryegrass, 386

CROFT, B. A. see COOP, L. B., 485

DASHIELL, K. E. see AKEM, C. N., 325
DASHIELL, K. E. AND AKEM, C. N. Yield losses in soybeans from frogeye leaf spot caused by *Cercospora sojina*, 465
DENNILL, G. B. A pruning technique for

saving vineyards severely infested by the grape vine bud mite *Colomerus* vitis (Pagenstecher) (Eriophyidae), 310

Despraux, D. see Gohet, E., 234 Dobson, H. M. see Symmons, P. M., 136 Dolbeer, R. A. see Mason, J. R., 294

Doster, M. A. and Fry, W. E. Evaluation by computer simulation of strategies to time metalaxyl applications for improved control of potato late blight, 209

DOYLE, C. J. Mathematical models in weed management, 432

Dreishpoun, J. see Shtienberg, D., 381 Droby, S. see Wilson, C. L., 172

EL MELEIGI, M. A. see EL-HAG, E. A. T., 65

EL MELEIGI, M. A. see EL-HAG, E. A. T., 70

EL-HAG, E. A. T. AND EL MELEIGI, M. A. Insect pests of spring wheat in Central Saudi Arabia, 65

EL-HAG, E. A. T. AND EL MELEIGI, M. A. Bionomics of the wheat leafminer Agromyza sp. (Diptera: Agromyzidae) in Central Saudi Arabia, 70

ELLIOT, P. C. AND MOODY, K. Influence of weed-control practices in the first crop on the tillage requirements for the succeeding crops in an upland rice-maize-cowpea cropping sequence. 28

ELLIOTT, R. H., MANN, L., SPURR, D. T. AND SACHER, E. R. Distribution of dye deposits from a small-plot track sprayer on biological and artificial targets, 129

EMMETT, B. J. see Young, A. G., 413 ERINLE, I. D. see CHINDO, P. S., 62

FENLON, J. S. see FERMOR, T. R., 271
FERMOR, T. R., HENRY, M. B., FENLON, J.
S., GLENISTER, M. J., LINCOLN, S. P.
AND LYNCH, J. M. Development and
application of a biocontrol system for
bacterial blotch of the cultivated
mushroom, 271

FISHLER, G. see BUTLER, G. D., 461 FITT, B. D. L. see BATEMAN, G. L., 479 FLYNN, J. L. see OGUNWOLU, E. O., 57

FOLWELL, R. J., COOK, R. J., HEIM, M. N. AND MOORE, D. L. Economic significance of take-all on winter wheat in the Pacific Northwest, USA, 391

FRIESEN, G. H. see WALL, D. A., 496 FRIESEN, G. H. AND WALL, D. A. Control of Sinapis arvensis in safflower with post-emergence herbicides, 74

FRY, W. E. see DOSTER, M. A., 209

GHATE, S. R., SUMNER, D. R. AND PHATAK, S. C. Stand and yield of cucumber seeded with gel and fungicides in various tillage systems, 23

GIGI, D. P., MUTEMEREWA, S., MOYO, G. AND NEELEY, D. Assessment and control of losses caused by insect pests in small farmers' stores in Zimbabwe, 287

GLEN, D. M., WILTSHIRE, C. W. AND BUTLER, R. C. Slug population changes following molluscicide treatment in relation to distance from edge of treated area. 408

GLENISTER, M. J. see FERMOR, T. R., 271 GOHET, E., VAN CANH, T., LOUANCHI, M. AND DESPRAUX, D. New developments in chemical control of white root disease of *Hevea brasiliensis* in Africa, 234

GOLD, C. S., ALTIERI, M. A. AND BELL-OTTI, A. C. Survivorship of the cassava whiteflies Aleurotrachelus socialis and Trialeurodes variabilis (Homoptera: Aleyrodidae) under different cropping systems in Colombia, 305

GOPALAN, M. see SENGUTTUVAN, T., 125 GREEN, D. I. see YOUNG, A. G., 413

GRIEFITHS, D. C., MANIAR, S. P., MER-RITT, L. A., MUDD, A., PICKETT, J. A., PYE, B. J., SMART, L. E. AND WAD-HAMS, L. J. Laboratory evaluation of pest management strategies combining antifeedants with insect growth regulator insecticides, 145

GUERRA-SOBREVILLA, L. Parasitoids of the grapeleaf skeletonizer, Harrisina brillians Barnes and McDunnough (Lepidoptera: Zygaenidae) in north-

western Mexico, 501

Guha, P. Control of *Chara* with oxadiazon and copper sulphate in water-logged rice fields in India, 371

HAGGAR, R. J., CLEAR HILL, B. H., ISAAC, S. P. AND STANDELL, C. J. Regulating sward growth and animal output with mefluidide, 178

 HALL, F. R. see ADAMS, A. J., 106
 HALL, J. S. AND HARMAN, G. E. Efficacy of oil treatments of legume seeds for control of Aspergillus and Zabrotes,

HALL, J. S. AND HARMAN, G. E. Protection of stored legume seeds against attack by storage fungi and weevils: mechanism of action of lipoidal and oil seed treatments, 375

HARMAN, G. E. see HALL, J. S., 315 HARMAN, G. E. see HALL, J. S., 375 HARMAN, G. E. Seed treatments for his

Harman, G. E. Seed treatments for biological control of plant disease, 166
Hasanuddin, A. see Sama, S., 34
Heim, M. N. see Folwell, R. J., 391
Helson, B. V. see Payne, N. J., 357
Henry, M. B. see Fermor, T. R., 271
Hensley, S. D. see Ogunwolu, E. O., 57
Hibino, H. see Sama, S., 34
Hodoson, C. J. see Afun, J. V. K., 363
Horne, J. and Balley, P. Bruchus

pisorum L. (Coleoptera, Bruchidae) control by a knockdown pyrethroid in field peas, 53

Imbe, T. see Shahjahan, M., 195 Isaac, S. P. see Haggar, R. J., 178

Jackai, L. E. N., Laboratory and screenhouse assays for evaluating cowpea resistance to the legume pod borer, 48 Jackai, L. E. N., see Afun, J. V. K., 363 Jalani, B. S. see Shahjahan, M., 195

JOHNSON, K. B. AND RADCLIFFE, E. B. Validation of a model simulating the feeding effects of the potato leafhopper (*Empoasca fabae*) on potato growth, 416

JOHNSTONE, D. R. Variations in insecticide dose received by settled locusts: a computer model for ultra-lowvolume spraying, 183

KABISSA, J. C. B. Studies on the performance of the 'Electrodyn' on cotton in eastern Tanzania, 354

KARJALAINEN, R. see PELTONEN, J., 260

KARJALAINEN, R. see PELTONEN, J., 200 KEARNEY, P. C. see MULBRY, W., 334 KHAN, F. A. see CHINDO, P. S., 62 KITTILA, S. see PELTONEN, J., 260 KOK, L. T. see VAIL, K. M., 199 KUC, J. see MUCHARROMAH, E., 265

Lambkin, T. M. see Collins, P. J., 215
Lemanceau, P. and Alabouvette, C.
Biological control of fusarium disease by fluorescent *Pseudomonas* and non-pathogenic *Fusarium*, 279
Lewis, G. C. see Cook, R., 403

Lewis, J. A. AND PAPAVIZAS, G. C. Biocontrol of plant diseases: the approach for tomorrow, 95

Lewis, J. A. AND PAPAVIZAS, G. C. Biocontrol of cotton damping-off caused by *Rhizoctonia solani* in the field with formulations of *Trichoderma* spp. and *Gliocladium virens*, 396

Lincoln, S. P. see Fermor, T. R., 271 Lindquist, R. K. see Adams, A. J., 106 Linz, G. see Mason, J. R., 294 Louanchi, M. see Gohet, E., 234 Lynch, J. M. see Fermor, T. R., 271

MACDONALD, O. C. Responses of the alien leaf miners Liriomyza trifolii and Liriomyza huidobrensis (Diptera: Agromyzidae) to some pesticides scheduled for their control in the UK, 509

Maniar, S. P. see Griffiths, D. C., 145 Mann, L. see Elliott, R. H., 129 Manwan, I. see Sama, S., 34

MARCUS, R. Deterministic and stochastic logistic models for describing increase of plant diseases, 155

Mason, J. R., Nuechterlein, G., Linz, G., Dolbeer, R. A. and Otis, D. L.

Oil concentration differences among sunflower achenes and feeding preferences of red-winged blackbirds, 294

MATTHEWS, G. A. see OMAR, D., 41
MAUDE, R. B. see PETCH, G. M., 117
MCAVOY, T. J. see VAIL, K. M., 199
MCGILL, N. G. see ALLSOPP, P. G., 205
MCLAUGHLIN, R. see WILSON, C. L., 172
MENN, J. J. Prospects and status for development of novel chemicals for IPM in cotton, 347

MERRITT, L. A. see GRIFFITHS, D. C., 145 MILLER, P. C. H. see THOMPSON, J. F., 254 MILLER, S. F. see COOP, L. B., 485 MIZEN, K. A. see COOK, R., 403 MOODY, K. see ELLIOT, P. C., 28 MOODY, K. see PABLICO, P., 45

Moore, D. L. see Folwell, R. J., 391 Moyo, G. see Giga, D. P., 287

MUCHARROMAH, E. AND KUC, J. Oxalate and phosphates induce systemic resistance against diseases caused by fungi, bacteria and viruses in cucumber, 265

MUDD, A. see GRIFFITHS, D. C., 145
MULBRY, W. AND KEARNEY, P. C. Degradation of pesticides by microorganisms and the potential for genetic manipulation, 334

Murphy, C. F. see Coop, L. B., 485 Murray, P. J. see Clements, R. O., 152 Mutemerewa, S. see Giga, D. P., 287

Neeley, D. see Giga, D. P., 287
Nelms, C. O. see Brugger, K. E., 455
Nuechterlein, G. see Mason, J. R., 294
Nyirenda, G. K. C. Effect of swath
width, time of application and height
on the efficacy of very-low-volume
(VLV) water-based insecticides on
cotton in Malawi, 111

Ogunwolu, E. O., Reagan, T. E., Flynn, J. L. and Hensley, S. D. Effects of *Diatraea saccharalis* (F.) (Lepidoptera: Pyralidae) damage and stalk rot fungi on sugar-cane yield in Louisiana, 57

OMAR, D. AND MATTHEWS, G. A. Influence of formulation and spray droplet size upon the persistence of permethrin deposits on brussels sprout leaves, 41

Omotade, M. A. see Babatola, J. O., 121 Omura, T. see Shahjahan, M., 195 Otis, D. L. see Mason, J. R., 294

Pablico, P. and Moody, K. Effect of fentin acetate on wet-seeded rice, *Pistia stratiotes* and *Azolla pinnata*, 45 Papavizas, G. C. see Lewis, J. A., 95 Papavizas, G. C. see Lewis, J. A., 396 Parker, C. Protection of crops against parasitic weeds, 6

PAYNE, N. J., SUNDARAM, K. M. AND HELSON, B. V. Airborne permethrin and off-target deposits from an aerial ultra-low-volume silvicultural spray, 357

- Peltonen, J., KITTILA, S., Peltonen-Sainio, P. and Karjalainen, R. Use of foliar-applied urea to inhibit the development of *Septoria nodorum* in spring wheat, 260
- PELTONEN-SAINIO, P. see PELTONEN, J., 260
- Petch, G. M., Maude, R. B. and White, J. G. Effect of film-coat layering of metalaxyl on the germination of carrot seeds, their emergence and the control of cavity spot, 117
- PHATAK, S. C. see GHATE, S. R., 23
 PICKETT, J. A. see GRIFFITHS, D. C., 145
 PIMENTEL, D. Diversification of biological control strategies in agriculture,
- POLLEY, R. W. see COOK, R. J., 504 PORT, G. R. see YOUNG, A. G., 413 PYE, B. J. see GRIFFITHS, D. C., 145
- QUIROGA, D., ARRETZ, P. AND ARAYA, J. E. Sucking insects damaging jojoba, Simmondsia chinensis (Link) Schneider, and their natural enemies, in the North Central and Central regions of Chile, 469
- RADCLIFFE, E. B. see JOHNSON, K. B., 416
 REAGAN, T. E. see OGUNWOLU, E. O., 57
 ROBERTSON, L. N. Soil-insecticide bioassays and seed treatments for soilinsect control in Central Queensland,
 Australia, 293
- ROELFS, A. P. see SECK, M., 222 ROGERS, D. J. see BRIER, H. B., 320

- SACHER, E. R. see ELLIOTT, R. H., 129
 SAMA, S., HASANUDDIN, A., MANWAN, I.,
 CABUNAGAN, R. C. AND HIBINO, H.
 Integrated management of rice
 tungro disease in South Sulawesi,
 Indonesia, 34
- Seck, M., Roelfs, A. P. and Teng, P. S. Influence of leaf position on yield loss caused by wheat leaf rust in single tillers, 222
- SEE, K. A. see Brown, J. K. M., 445
 SENGUTTUVAN, T., GOPALAN, M. AND
 CHELLIAH, S. Impact of resistance
 mechanisms in rice against the brown
 planthopper, Nilaparvata lugens Stål
- (Homoptera: Delphacidae), 125
 SHAHJAHAN, M., ZAKRI, A. H., IMBE, T.,
 JALANI, B. S. AND OMURA, T. Efficient
 method of determining tungro virus
 resistance in rice (Oryza sativa L.),
 195
- SHTIENBERG, D. AND DREISHPOUN, J. Suppression of Alternaria leaf spot in Pima cotton by systemic fungicides, 381
- SISSOKO, M. see SYMMONS, P. M., 136 SLATER, S. E. see BROWN, J. K. M., 445 SMART, L. E. see GRIFFITHS, D. C., 145 SMITH, B. D. see COUCH, H. B., 386 SNEH, B. see BROZA, M., 229 SPURR, D. T. see ELLIOTT, R. H., 129
- SNEH, B. see BROZA, M., 229 SPURR, D. T. see ELLIOTT, R. H., 129 STAFFORD, J. V. see THOMPSON, J. F., 254 STANDELL, C. J. see HAGGAR, R. J., 178 SUMNER, D. R. see GHATE, S. R., 23
- SUNDARAM, K. M. see PAYNE, N. J., 357
 SUTHERST, R. W. Predicting the survival
 of immigrant insect pests in new
 environments, 331
- Symmons, P. M., Dobson, H. M. And Sissoko, M. Pesticide droplet size and efficacy: a series of trials on grasshoppers, 136

- TENG, P. S. see SECK, M., 222
 THOMAS, M. R. see COOK, R. J., 504
 THOMPSON, J. F., STAFFORD, J. V. AND
 MILLER P. C. Potential for automatic
- THOMPSON, J. F., STAFFORD, J. V. AND MILLER, P. C. Potential for automatic weed detection and selective herbicide application, 254
 TREVERROW, N. L. see COLLINS, P. J., 215
 - 1 REVERROW, N. L. see COLLINS, P. J., 215
- VAIL, K. M., KOK, L. T. AND MCAVOY, T. J. Cultivar preferences of lepidopterous pests of broccoli, 199 VAN CANH, T. see GOHET, E., 234
- WADHAMS, L. J. see GRIFFITHS, D. C., 145 WALL, D. A. see FRIESEN, G.H., 74
- WALL, D. A. AND FRIESEN, G. H. Effectiveness of selective herbicides for the control of annual weed species in lathyrus (*Lathyrus sativus L.*,), 496
- WHITE, J. G. see PETCH, G. M., 117
 WILSON, C. L., WISNIEWSKI, M. E., BILES,
 C. L., MCLAUGHLIN, R., CHALUTZ,
 E. AND DROBY, S. Biological control
 of post-harvest diseases of fruits and
 vegetables: alternatives to synthetic
 fungicides. 172
- WILSON, F. D. see BUTLER, G. D., 461 WILTSHIRE, C. W. see GLEN, D. M., 408 WISNIEWSKI, M. E. see WILSON, C. L., 172
- Young, A. G., Port, G. R., Emmett, B. J. and Green, D. I. Development of a forecast of slug activity: models to relate slug activity to meteorological conditions, 413
- ZAKRI, A. H. see SHAHJAHAN, M., 195